

Analysis of an infinite sequence

Assignment

Now we will expand our investigation to determine the sum of the infinite sequence t_n , where

$$t_0=1, t_1=(x \ln a) 1, t_2=(x \ln a) 2 2 x 1, t_3=(x \ln a) 3 3 x 2 x 1, \dots$$

Define $T_n(a,x)$ as the sum of the first n terms, for various values of a and x .

e.g. $T_9(2,9)$ is the sum of the first nine terms when $a=2$ and $x=5$.

Let $a=2$. Calculate $T_9(2,x)$ for various positive values of x . Using technology, plot the relation between $T_9(2,x)$ and x . Describe what you notice from your plot.

Let $a=3$. Calculate $T_9(3,x)$ for various positive values of x . Using technology, plot the relation between $T_9(3,x)$ and x . Describe what you notice from your plot.

Continue with this analysis to find the general statement for $T_n(a,x)$ as n approaches ∞ .

Test the validity of the general statement with other values of a and x .

Discuss the scope and/or limitations of the general statement.

Explain how you arrived at the general statement.

Work

1. First we will use the mathematical program DeriveTM 6 to calculate the sum of the first 9 terms of the sequence when $a=2$, for various values of x .

The expression has the following form:

SUM(Expression, summation parameter, lower boundary, upper boundary)

Which in our case makes:

SUM(((x*ln(2))^n)/n!,n,0,9)

DeriveTM 6 then shows the following:

$$n=9(x \cdot \ln(2))^n n!$$

Option Simplify gives us the expanded form of the sum:

$$x^9 \ln(a)^9 362880 + x^8 \ln(a)^8 40320 + x^7 \ln(a)^7 5040 + x^6 \ln(a)^6 720 + x^5 \ln(a)^5 120 + x^4 \ln(a)^4 24 + x^3 \ln(a)^3 6 + x^2 \ln(a)^2 2 + x \cdot \ln(a) + 1$$

By using the button \approx , the following results were reached for arbitrarily chosen values of x:

n	a	x	T _n (a,x)
9	2	0.1	1.071773462
9	2	0.2	1.148698354
9	2	0.5	1.414213562
9	2	1	2
9	2	2	3.999991747
9	2	3	7.999488457
9	2	8	241.6244730
9	2	10	857.4734736

TABLE 1. Calculated sums of 9 terms of the sequence for various values of x when a=2

The calculated value for x=1 stands out, being an integer.

We now write a MATLAB[©] program to plot the sum as a function of x.

The main program code:

```
a=2;
n=9; % the number of terms contained in the sum
L=log(a); % To avoid needless calculation in each iteration,
% the natural logarithm is calculated at the beginning.
Tn=0; % Setting the initial sum of the sequence to zero.
```

---- OSTATAK TEKSTA NIJE PRIKAZAN. CEO RAD MOŽETE PREUZETI NA SAJTU
WWW.MATURSKI.NET ----

[BESPLATNI GOTOVI SEMINARSKI, DIPLOMSKI I MATURSKI TEKST](#)


RAZMENA LINKOVA - RAZMENA RADOVA

RADOVI IZ SVIH OBLASTI, POWERPOINT PREZENTACIJE I DRUGI EDUKATIVNI MATERIJALI.

WWW.SEMINARSKIRAD.ORG

WWW.MAGISTARSKI.COM

WWW.MATURSKIRADOVI.NET



NA NAŠIM SAJTOVIMA MOŽETE PRONAĆI SVE, BILO DA JE TO [SEMINARSKI, DIPLOMSKI](#) ILI [MATURSKI](#) RAD, POWERPOINT PREZENTACIJA I DRUGI EDUKATIVNI MATERIJAL. ZA RAZLIKU OD OSTALIH MI VAM PRUŽAMO DA POGLEDATE SVAKI RAD, NJEGOV SADRŽAJ I PRVE TRI STRANE TAKO DA MOŽETE TAČNO DA ODABERETE ONO ŠTO VAM U POTPUNOSTI ODGOVARA. U BAZI SE NALAZE [GOTOVI SEMINARSKI, DIPLOMSKI I MATURSKI RADOVI](#) KOJE MOŽETE SKINUTI I UZ NJIHOVU POMOĆ NAPRAVITI JEDINSTVEN I UNIKATAN RAD. AKO U [BAZI](#) NE NAĐETE RAD KOJI VAM JE POTREBAN, U SVAKOM MOMENTU MOŽETE NARUČITI DA VAM SE IZRADI NOVI, UNIKATAN SEMINARSKI ILI NEKI DRUGI RAD RAD NA LINKU [IZRADA RADOVA](#). PITANJA I ODGOVORE MOŽETE DOBITI NA NAŠEM [FORUMU](#) ILI NA

maturskiradovi.net@gmail.com

